| APPENDIX C /HR Modeling Run Results for Shaded Fuelbreak Scenar |
|--|
| |
| |
| /HP Modeling Pun Results for Shadad Euclbrook Soons |
| |
| The modeling Run Results for Shaded Fuelbreak Scenar |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

Weighted Habitat Value Comparison Report

| FIRST CONDITION SELECTION CRITERIA | SECOND CONDITION SELECTION CRITERIA |
|--|-------------------------------------|
| Locations: | Locations: |
| County: BUTTE | County: BUTTE |
| County. BOTTE | County. BollB |
| Habitats/Seral Stages: | Habitats/Seral Stages: |
| Type/Stage 1: BOP3P | Type/Stage 1: BAR1 |
| Type/Stage 2: BOP3M | Type/Stage 2: URB1 |
| Type/Stage 3: BOP3D | Type/Stage 3: AGS2D |
| Type/Stage 4: VRI2M | Type/Stage 4: MCH3S |
| Type/Stage 5: VRI2D | Type/Stage 5: MCH3M |
| Type/Stage 6: MHW3D | Type/Stage 6: BOP3P |
| Type/Stage 7: MCH3M | Type/Stage 7: BOP3M |
| Type/Stage 8: AGS2D | Type/Stage 8: BOP3D |
| Type/Stage 9: FEW2D | Type/Stage 9: VRI2P |
| Type/Stage 10: URB1 | Type/Stage 10: VRI2M |
| Type/Stage 11: BAR1 | Type/Stage 11: MHW3M |
| Type/Stage 11. D/M(1 | Type/Stage 12: FEW2D |
| Excluded Elements: | Type/Stage 13: BOP4P |
| KELP | Type/Stage 14: MHW4P |
| KELF | Type/Stage 14. WITW4F |
| Element Levels (E,S,P): | Excluded Elements: |
| REPRO: E | KELP |
| COVER: E | KELI |
| FEEDING: E | Element Levels (E,S,P): |
| reeding. e | REPRO: E |
| . Unbitet Suitability Levels (U.M.L.): | COVER: E |
| Habitat Suitability Levels (H,M,L): REPRO: L | FEEDING: E |
| COVER: L | PEEDING. E |
| FEEDING: L | Habitat Suitability Levels (H,M,L): |
| FEEDING. L | REPRO: L |
| Weights: | COVER: L |
| BLUE OAK-FOOTHILL PINE3P 19 | FEEDING: L |
| BLUE OAK-FOOTHILL PINE3M 24 | reeping. L |
| BLUE OAK-FOOTHILL PINE3D 46 | Weights: |
| VALLEY FOOTHILL RIPARIAN2M 1 | BARRENNo 1 |
| VALLEY FOOTHILL RIPARIAN2D 1 | URBANNo 7 |
| MONTANE HARDWOOD3D 1 | ANNUAL GRASS2D 1 |
| MIXED CHAPARRAL3M 1 | MIXED CHAPARRAL3S 1 |
| ANNUAL GRASS2D 1 | MIXED CHAPARRAL3M 1 |
| FRESH EMERGENT WETLAND2D 1 | BLUE OAK-FOOTHILL PINE3P 19 |
| URBANNo 7 | BLUE OAK-FOOTHILL PINE3M 23 |
| BARRENNo 1 | BLUE OAK-FOOTHILL PINE3D 45 |
| DARKENNO | VALLEY FOOTHILL RIPARIAN2P 1 |
| • | VALLEY FOOTHILL RIPARIAN2M 1 |
| | MONTANE HARDWOOD3M 1 |
| | FRESH EMERGENT WETLAND2D 1 |
| | BLUE OAK-FOOTHILL PINE4P 2 |
| | MONTANE HARDWOOD4P 1 |
| | |

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

The results of this report are meant to indicate trends only. The numerical calculations are not intended to represent any generally accepted method of habitat evaluation (e.g. HEP). Predictions should be validated with field surveys by a professional wildlife biologist.

| ID Species Name | Difference |
|---|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B326 ASH-THROATED FLYCATCHER | 2.80 |
| *9 *7 *3 0 0 *7 *7 | 48.90 |
| *7 *7 *9 *7 *3 0 0 *7 *10 *10 | 51.70 |
| R051 RACER | 2.80 |
| *7 *7 *7 *3 *3 *10 *7 | 31.30 |
| *10 *7 *3 *7 *7 *7 *7 *3 *7 *7 *7 | 34.10 |
| B265 GREAT HORNED OWL | 3.00 |
| 0 0 0 0 0 0 0 0 *10 *10 | 8.00 |
| *10 *10 0 0 0 0 0 0 0 0 0 *10 *10 | 11.00 |
| B408 PHAINOPEPLA | 3.00 |
| *9 *3 *3 0 *10 | 24.10 |
| *10 0 0 *9 *3 *3 *3 *10 *7 | 27.10 |
| R057 GOPHER SNAKE | 3.00 |
| *10 *10 *3 *10 *7 *7 *10 *3 *3 | 64,20 |
| *3 *10 *10 *7 *10 *10 *3 *10 *10 *3 *3 *10 *7 | 67.20 |
| B287 ANNA'S HUMMINGBIRD | 3.10 |
| *10 *7 *3 *5 *3 *10 *10 | 59.10 |
| *10 *10 *10 *10 *7 *3 *6 *5 *3 *10 *7 | 62.20 |
| B344 BARN SWALLOW | 3.10 |
| *10 *7 *3 *7 *3 *7 *10 *10 0 | 54.00 |
| 0 *10 *7 *7 *10 *7 *3 *7 *7 *5 *10 *10 *7 | 57.10 |
| M051 BLACK-TAILED JACKRABBIT | 3.10 |
| *7 *3 *3 *5 0 *3 | 23.80 |
| *3 0 *10 *5 *7 *3 *3 *4 *7 *7 | 25.80 |
| B512 DARK-EYED JUNCO | 3.20 |
| *10 *7 *4 *7 *4 *4 *3 *7 | |
| *7 *3 *3 *10 *7 *4 *10 *7 *7 *10 *10 | 60.70 |
| B277 COMMON POORWILL | 63.90 |
| *10 *5 0 0 *7 0 *10 | |
| *10 0 *10 *7 *10 *5 0 *4 *10 *7 | 32.20 |
| B354 COMMON RAVEN | 35.80 |
| | 3.60 |
| | 52.70 |
| *9 0 *9 *9 *10 *7 *3 *9 *3 *3 0 *10 *10 R048 RINGNECK SNAKE | 56.30 |
| | 3.70 |
| *10 *3 *7 *3 *10 *3 *3 *3 | 31.80 |
| *3 *3 *10 *10 *10 *3 *7 *7 *3 *3 *10 *3 | 35.50 |
| M119 BRUSH MOUSE | 3.80 |
| *7 *3 *7 *3 *10 *3 | 22.70 |
| *3 *10 *10 *7 *3 *7 *7 *7 *7 | 26.50 |
| B262 BARN OWL | 4.10 |
| *10 *7 *7 *7 *7 0 0 *10 *0 | 44.70 |
| *0 *10 0 *9 *7 *10 *7 *9 *7 *7 0 *10 *10 Total Number of Species Afforded: 354 | 48.80 |

Total Number of Species Affected: 354

How the seral stage values are calculated:

Each preference value is translated from the letter codes H, M, L and blank to numeric codes 1, 0.66, 0.33 and 0 respectively. The preference values for reproduction, feeding and cover are averaged using the method you selected during the query process (geometric or arithmetic). For display purposes the resulting mean is multiplied by 10 to produce a value between 1 and 10 as the overall rating for the habitat/seral stage combination. When a stage is not rated for a species, it will not be displayed on this report As a result, the number of stage ratings may not equal the number of stages selected in the query process. An asterisk denotes a reproductive value is included in the habitat/seral stage rating. See the CWHR documentation for further information.

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|--|------------------------------------|
| Condition 1 Habitat Suitability Values | Condition 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B274 NORTHERN SAW-WHET OWL | 2.30 |
| 0 0 0 0 0 0 *3 | 2.30 |
| *3 0 0 0 0 0 0 0 *8 *8 | 4.60 |
| B341 NORTHERN ROUGH-WINGED SWALLOW | 2.30 |
| *10 *9 *7 *9 *7 *7 *3 0 0 *10 *7 | 81.90 |
| *7 *10 0 *3 *3 *10 *9 *7 *10 *9 *9 0 *10 *10 | 84.20 |
| B410 LOGGERHEAD SHRIKE | 2.30 |
| *10 *8 *3 *3 0 0 | 37.80 |
| 0 0 *3 *3 *10 *8 *4 *3 *10 *3 | 40.10 |
| B538 HOUSE FINCH | 2.30 |
| *10 *7 *3 *7 *3 0 *3 *10 | 58.30 |
| *10 *5 *3 *10 *7 *3 *10 *7 0 *10 0 | 60.60 |
| M149 GRAY FOX | 2.30 |
| *10 *9 *5 *7 *5 *5 *9 *4 *5 *3 | 70.10 |
| *3 *4 *5 *9 *10 *9 *5 *5 *7 *9 *5 *10 *8 | 72.40 |
| R022 WESTERN FENCE LIZARD | 2.30 |
| *10 *10 *7 *10 *7 *3 *7 *3 *3 | 78.70 |
| *3 *3 *10 *7 *10 *10 *7 *10 *10 *3 *10 *7 | 81.00 |
| R062 WESTERN TERRESTRIAL GARTER SNA | 2.30 |
| *10 *10 *7 *10 *8 *3 *7 *10 | 77.20 |
| *7 *7 *10 *10 *7 *10 *10 *7 *10 *10 *7 | 79.50 |
| B389 AMERICAN ROBIN | 2,40 |
| *9 *3 *0 *3 *3 *7 0 0 *10 | 32.70 |
| *10 0 0 0 *9 *3 *0 *5 *3 *5 *9 *10 | 35.10 |
| B532 BULLOCK'S ORIOLE | 2.40 |
| *5 *4 0 0 0 *4 *10 | 26.60 |
| *10 *5 *4 0 0 0 *4 *10 *8 | 29.00 |
| M127 DUSKY-FOOTED WOODRAT | 2.40 |
| *7 *10 *3 *7 *3 *7 *10 | 54.40 |
| *7 *10 *7 *10 *3 *7 *7 *10 *7 *10 | 56.80 |
| M162 STRIPED SKUNK | 2.40 |
| *10 *7 *3 *9 *5 *3 *9 *8 *8 *5 | 57,90 |
| *5 *8 *8 *9 *10 *7 *3 *10 *9 *7 *8 *5 *5 | 60.30 |
| R036 WESTERN SKINK | 2.40 |
| *10 *10 *7 *10 *7 *3 *7 *3 | 76.40 |
| *3 *7 *7 *10 *10 *7 *10 *10 *7 *10 *7 | 78.80 |
| R049 SHARP-TAILED SNAKE | 2.40 |
| *10 *10 *7 *7 *7 *3 *10 | 76.10 |
| *10 *10 *10 *10 *7 *7 *7 *7 *10 *7 | 78.50 |
| B415 CASSIN'S VIREO | 2.50 |
| *6 *5 *3 *7 *3 *4 0 | 40.70 |
| 0 *6 *5 *3 *8 *7 *10 *6 *10 | 43.20 |
| M072 CALIFORNIA GROUND SQUIRREL | 2.50 |
| *7 *3 *3 *3 *3 *3 *10 *10 | 44.80 |
| *10 *10 *7 *3 *7 *3 *3 *7 *3 *7 *7 | 47.30 |
| B543 LESSER GOLDFINCH | 2.60 |
| *8 *7 *3 *4 *3 *4 *3 0 *10 | 53.80 |
| *10 0 *7 *3 *8 *7 *3 *8 *4 *4 *8 *8 | 56.40 |
| B108 TURKEY VULTURE | 2.70 |
| *10 *9 *7 *9 *7 *7 *10 *5 *9 | 79.90 |
| *9 *5 *10 *10 *7 *10 *9 *7 *10 *9 | 79.90 82.60 |
| | 02.UV |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|--|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| M081 BOTTA'S POCKET GOPHER | 2.00 |
| *3 *10 *7 *3 | 12.20 |
| 3 *7 *10 *3 *3 *7 *3 *3 | 14.20 |
| 3255 MOURNING DOVE | 2.10 |
| 8 *10 *7 *4 *3 *7 *4 *10 | 77.60 |
| *10 *4 *4 *8 *10 *7 *4 *4 *10 *10 *10 | 79.70 |
| 3282 WHITE-THROATED SWIFT | 2.10 |
| 10 *10 *10 *7 *7 *10 *9 0 0 *5 *10 | 96.70 |
| *10 *5 0 *9 *9 *10 *10 *10 *9 *7 *10 0 *10 *10 | 98.80 |
| 3377 BLUE-GRAY GNATCATCHER | 2.10 |
| 10 *8 *7 *3 *3 *7 | 68.90 |
| 3 *7 *10 *8 *7 *3 *3 *5 *10 *7 | 71.00 |
| 3489 CHIPPING SPARROW | 2.10 |
| 10 *7 *7 0 *3 0 | 65.50 |
| 0 0 *10 *7 *7 0 0 *7 *10 *10 | 67.60 |
| 3544 LAWRENCE'S GOLDFINCH | 2.10 |
| 8 *4 *4 *3 *4 *3 0 *4 | 28.80 |
| 4 0 *3 *3 *8 *4 *5 *4 *4 *8 *5 | 30.90 |
| MO34 HOARY BAT | 2.10 |
| 7 *7 *5 0 0 *5 0 0 0 0 | 53.20 |
| 0 0 0 *7 *7 *5 0 0 *7 0 *10 *10 | 55.30 |
| 4075 GOLDEN-MANTLED GROUND SQUIRREL | 2.10 |
| 5 *5 *3 *7 0 | 2.00 |
|) *8 *7 *4 *5 *7 *10 | 4.10 |
| M181 MULE DEER | 2.10 |
| 9 *9 *5 *7 *4 *5 *9 *4 0 *4 | 65.30 |
| 4 *4 *5 *9 *9 *9 *5 *9 *7 *8 0 *8 *7 | 67.40 |
| 2029 COAST HORNED LIZARD | 2.10 |
| 7 *3 *3 *3 | 21.00 |
| 10 *3 *7 *3 *7 *3 *3 | 23.10 |
| 1061 COMMON GARTER SNAKE | 2.10 |
| 7 *7 *7 *10 *10 *3 *10 *10 *10 | 64.00 |
| 10 *10 *10 *7 *7 *7 *10 *10 *7 *10 *7 *7 | 66.10 |
| 1076 WESTERN RATTLESNAKE | 2.10 |
| 10 *10 *10 *10 *3 *10 *7 *3 *3 *3 | 94.90 |
| 3 *3 *7 *10 *10 *10 *10 *10 *10 *7 *3 *10 *7 | 97.00 |
| 123 RED-TAILED HAWK | 2.20 |
| 8 *7 *5 *4 *3 *5 0 *5 0 *10 *7 | 64.00 |
| 7 *10 *5 0 0 *8 *7 *5 *4 *4 *7 0 *10 *10 | 66.20 |
| 343 CLIFF SWALLOW | 2.20 |
| 10 *5 *0 *10 *10 *7 *8 *8 *10 | 43.00 |
| 10 *8 *7 *7 *10 *5 *0 *10 *10 *8 *10 | 45.00 |
| 117 DEER MOUSE | 43.20 |
| 7 *7 *3 *7 *3 *3 *10 *7 *7 *3 | 52.10 |
| 3 *7 *7 *7 *10 *7 *7 *3 *7 *7 *3 *7 *7 | |
| 127 AMERICAN KESTREL | 54.30 |
| | 2.30 |
| | 7.50 |
| *10 *5 0 0 0 0 0 0 0 0 *8 *8 140 CALIFORNIA QUAIL | 9.80 |
| 140 CALIFORNIA QUAIL 10 *10 *8 *10 *8 *5 *7 *8 *9 | 2.30 |
| | 87.90 |
| 9 *8 *10 *7 *10 *10 *8 *10 *10 *7 *10 *7 | 90.20 |

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|--|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B131 PRAIRIE FALCON | 1.80 |
| *9 *7 *7 *7 *7 *5 *10 0 0 | 68.40 |
| 0 0 *10 *5 *5 *9 *7 *7 *7 *7 0 *9 *9 | 70.20 |
| B141 MOUNTAIN QUAIL | 1.80 |
| *8 *8 *5 *6 *5 *8 0 *3 | 56,90 |
| *3 *4 0 *8 *8 *5 *6 *6 *8 *8 *10 | 58.70 |
| B311 WESTERN WOOD-PEWEE | 1.80 |
| *8 *8 *8 0 0 *8 *3 *6 | 72.90 |
| *6 *3 *3 *8 *8 *8 0 0 *8 *10 *10 | 74.70 |
| B528 BROWN-HEADED COWBIRD | 1.80 |
| *10 *7 *3 *4 *4 *0 *7 *5 *6 *10 | 59.60 |
| *10 *5 *7 *7 *10 *7 *3 *5 *4 *0 *6 *7 *5 | 61.40 |
| M001 VIRGINIA OPOSSUM | 1.80 |
| *7 *7 *3 *10 *10 *3 *3 *7 *7 | 51.70 |
| *7 *3 *7 *3 *7 *7 *3 *10 *10 *3 *7 *7 *7 | 53.50 |
| M113 WESTERN HARVEST MOUSE | 1.80 |
| *3 *3 *7 *3 *3 *10 *7 | 32.40 |
| *10 *7 *3 *3 *3 *3 *7 *7 *3 *7 *3 | 34.20 |
| M128 BUSHY-TAILED WOODRAT | 1.80 |
| *7 *3 *7 *7 *4 | 2.80 |
| *4 *7 *7 *7 *7 *7 | 4.60 |
| M152 RINGTAIL | 1.80 |
| *10 *7 *3 *9 *7 *3 *9 0 *0 | 52.80 |
| *0 0 *8 *9 *10 *7 *3 *10 *9 *3 *7 *3 | 54.60 |
| M153 RACCOON | 1.80 |
| *4 *5 *3 *5 *4 *3 *7 0 0 *7 | 42.40 |
| *7 0 *3 *7 *4 *5 *3 *5 *5 *5 0 *7 *7 | 44.20 |
| B286 BLACK-CHINNED HUMMINGBIRD | 1.90 |
| *8 *4 *4 *3 0 *10 | 32,20 |
| *10 0 0 *8 *4 *6 *4 *10 | 34.10 |
| M166 BOBCAT | 1.90 |
| *10 *7 *5 *10 *7 *5 *9 *4 *4 | 63.10 |
| *4 *5 *9 *10 *7 *5 *10 *10 *7 *4 *7 *7 | 65.00 |
| B267 NORTHERN PYGMY OWL | 2.00 |
| *6 *6 *5 0 0 *5 0 *10 | 57.70 |
| *10 0 0 *6 *6 *5 0 0 *6 *10 *10 | 59.70 |
| B296 ACORN WOODPECKER | 2.00 |
| *7 *6 *5 0 0 *5 *10 | 57.10 |
| *10 *7 *6 *5 0 0 *6 *10 *10 | 59.10 |
| B340 VIOLET-GREEN SWALLOW | 2.00 |
| *5 *5 *5 0 0 *5 0 0 0 *10 0 | 50.20 |
| 0 *10 0 0 0 *5 *5 *5 0 0 *5 0 *10 *10 | 52.20 |
| B417 HUTTON'S VIREO | 2.00 |
| *10 *9 *6 *6 *6 *5 *3 *7 | 74.30 |
| *7 *3 *3 *10 *9 *6 *8 *6 *5 *10 *10 | 76.30 |
| M028 CALIFORNIA MYOTIS | 2.00 |
| *7 *7 *5 *7 *5 *5 *7 0 0 *7 *0 | 59.70 |
| *0 *7 0 *7 *7 *7 *5 *7 *7 *7 0 *7 *7 | 61.70 |
| M032 BIG BROWN BAT | 2.00 |
| *10 *10 *10 *10 *10 *7 0 0 *10 *7 | 100.00 |
| *7 *10 0 *7 *7 *10 *10 *10 *10 *10 *10 0 *10 *10 | |
| , 20 0 , , 10 10 10 10 10 10 10 | 102.00 |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | | ference |
|---|------------------------------------|---------|
| Condition 1 Habitat Suitability Values | Condition 1 Sum of Weighted Values | |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values | .50 |
| 366 ROCK WREN | | 30 |
| 7 *7 *7 *7 *10 10 *7 *7 *7 *7 *7 *7 *7 *7 | 63.80 65.30 | |
| 10 *7 *7 *7 *7 *7 *7 *7 *7 *7 380 WESTERN BLUEBIRD | | 1.50 |
| | 67.90 | |
| 8 *7 *7 *3 0 *10 10 0 0 *8 *7 *7 0 *3 *10 *8 | 69.40 | |
| 10 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | | .50 |
| 10 *9 *6 *9 *7 *6 *10 *5 0 *5 0 | 74.60 | |
| *5 *5 *8 *10 *10 *9 *6 *10 *9 *9 0 *5 *5 | 76.10 | |
| M161 WESTERN SPOTTED SKUNK | | 1.50 |
| 10 *7 *3 *10 *7 *3 *10 *4 *3 | 55,70 | |
| 3 *4 *7 *10 *10 *7 *3 *10 *10 *7 *3 *3 | 57.20 | |
| .032 WESTERN TOAD | 1 | 1.60 |
| 7 *7 *3 *7 *7 *4 *3 *7 *7 *3 | 49.30 | |
| 3 *7 *5 *3 *7 *7 *3 *7 *7 *4 *7 *7 | 50.90 | |
| .046 BULLFROG | 1 | 1.60 |
| 7 *7 *7 *9 *7 *7 *7 *7 *9 *7 | 67.90 | |
| 7 *7 *7 *7 *7 *7 *7 *9 *9 *7 *9 *7 | 69.50 | |
| 3426 NASHVILLE WARBLER | 1 | 1.60 |
| 0 0 0 0 *5 *3 0 | 0.80 | |
| *3 *3 0 0 0 0 0 *8 0 *10 | 2.40 | |
| 477 LAZULI BUNTING | | 1.60 |
| 10 *7 *7 *10 *7 *3 *9 | 68.10 | |
| 3 *9 *10 *7 *7 *10 *10 *3 *10 *3 | 69.70 | |
| PLUMBEOUS VIREO | | 1.60 |
| 0 0 0 0 *4 0 | 0.40 | |
| 0 0 0 0 0 *10 0 *10 | 2.00 | 1.60 |
| 1047 DESERT COTTONTAIL | | 1.60 |
| 7 *7 *3 *7 *5 *9 *7 *10 | 53.30 | |
| 10 *7 *10 *9 *7 *7 *3 *7 *7 *7 | 34.90 | 1.60 |
| 2040 SOUTHERN ALLIGATOR LIZARD | | 1.60 |
| 10 *10 *7 *10 *10 *3 *10 *3 *3 | 79.30 | |
| 3 *3 *10 *10 *10 *10 *7 *10 *10 *3 *10 *3 3126 GOLDEN EAGLE | 80.90 | 1.70 |
| | 75.20 | 1.70 |
| 10 *9 *7 *3 *7 *7 *4 *5 0 0 *10 10 0 *5 *4 *4 *10 *9 *7 *3 *3 *9 0 *10 *10 | 75.20 | |
| 352 YELLOW-BILLED MAGPIE | | 1.70 |
| 0 0 0 0 *10 | 7.00 | |
| 10 0 0 0 0 0 *9 | 8.70 | |
| 3353 AMERICAN CROW | | 1.70 |
| 5 *3 *3 0 0 *3 0 *10 | 39.50 | |
| 10 0 *5 *3 *3 0 0 *3 *10 *3 | 41.20 | |
| 418 WARBLING VIREO | | 1.70 |
| 10 *10 *8 *8 *5 *5 *7 | 84.60 | |
| 7 *10 *10 *8 *8 *8 *7 *10 *10 | 86.30 | |
| 1487 RUFOUS-CROWNED SPARROW | | 1.70 |
| 7 *3 *3 *7 *10 | 37.30 | |
| 10 *10 *7 *7 *3 *3 0 *7 | 39.00 | |
| MO55 YELLOW-PINE CHIPMUNK | | 1.70 |
| 3 *7 | 1.00 | |
| 3 *7 *7 *10 | 2.70 | |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|---|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B307 NORTHERN FLICKER | 1.30 |
| *6 *6 *5 0 0 *5 0 0 *10 | 57.70 |
| *10 0 0 0 *6 *6 *5 0 0 *6 *7 *10 | 59.00 |
| B393 NORTHERN MOCKINGBIRD | 1.30 |
| *5 *5 *3 *3 *10 0 *10 | 44.10 |
| *10 0 *10 *10 *5 *5 *3 *3 *4 | 45.40 |
| B475 BLACK-HEADED GROSBEAK | 1.30 |
| *10 *10 *7 0 0 *10 *3 *10 | 81.70 |
| *10 0 *3 *10 *10 *7 0 0 *10 *10 *10 | 83.00 |
| M018 BROAD-FOOTED MOLE | 1.30 |
| *3 *10 *7 *10 | 9.00 |
| *10 *3 *10 *10 *3 *3 | 10.30 |
| M037 TOWNSEND'S BIG-EARED BAT | 1.30 |
| 0 0 0 0 0 0 0 0 *3 *0 | 2.30 |
| *0 *3 0 0 0 0 0 0 0 0 0 *7 0 | 3.60 |
| R042 NORTHERN ALLIGATOR LIZARD | 1.30 |
| *7 *3 | 1.00 |
| *3 *10 *10 | 2.30 |
| R058 COMMON KINGSNAKE | 1.30 |
| *7 *7 *10 *10 *3 *10 *7 *7 *3 | 65.70 |
| *3 *7 *10 *10 *7 *7 *7 *10 *10 *3 *7 *7 *3 | 67.00 |
| A039 PACIFIC CHORUS FROG | 1.40 |
| *7 *7 *10 *10 *7 *7 *10 *10 *5 | 67.80 |
| *5 *10 *7 *7 *7 *7 *10 *10 *7 *10 *7 *7 | 69.20 |
| B129 PEREGRINE FALCON | 1.40 |
| *9 *9 *9 *9 *9 *5 *10 0 *10 *10 | 89.60 |
| *10 *10 *10 *5 *5 *9 *9 *9 *9 *9 0 *9 *9 | 91.00 |
| B264 WESTERN SCREECH OWL | 1.40 |
| *10 *10 *9 0 0 *8 0 0 *10 | 90.80 |
| *10 0 0 0 *10 *10 *9 0 0 *10 *10 *10 *10 | 92.20 |
| B275 LESSER NIGHTHAWK | |
| *5 0 0 0 0 0 0 *10 | 10.10 11.50 |
| *10 0 *5 0 0 *4 0 0 *5 B358 OAK TITMOUSE | 11.30 |
| *8 *8 *4 *4 *8 *7 *10 | 76.90 |
| *10 *3 *7 *8 *8 *8 *3 *4 *8 *10 *7 | 78.30 |
| B362 WHITE-BREASTED NUTHATCH | 1.40 |
| *6 *6 *6 0 0 *6 *10 | 61.00 |
| *10 *6 *6 *6 0 0 *6 *9 *9 | 62.40 |
| B436 BLACK-THROATED GRAY WARBLER | 1.40 |
| *10 *10 *8 0 0 *8 0 0 | 78.80 |
| 0 0 0 *10 *10 *8 0 0 *10 *10 *10 | 80.20 |
| B505 SONG SPARROW | 1.40 |
| *3 *7 *3 *3 0 *10 *7 | 13,20 |
| *7 0 0 *3 *10 *7 *10 *3 *3 | 14.60 |
| M157 LONG-TAILED WEASEL | 1.40 |
| *10 *7 *5 *9 *7 *5 *4 *8 *3 | 64.80 |
| *3 *8 *8 *4 *10 *7 *5 *10 *9 *7 *4 *4 | 66.20 |
| B303 DOWNY WOODPECKER | 1.50 |
| *3 *3 *3 0 0 0 0 0 *10 | 36.40 |
| *10 0 0 0 *3 *3 *3 0 0 *7 *3 *7 | 37.90 |
| | |

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name Condition 1 Habitat Suitability Values | Difference |
|--|--|
| Condition 1 Habitat Suitability Values Condition 2 Habitat Suitability Values | Condition 1 Sum of Weighted Values Condition 2 Sum of Weighted Values |
| R059 CALIFORNIA MOUNTAIN KINGSNAKE | 1.00 |
| *3 *4 *4 *3 *3 *7 *7 *3 | 38.00 |
| *3 *3 *7 *3 *4 *4 *3 *3 *7 *4 *7 | 39.00 |
| R071 NIGHT SNAKE | 1.00 |
| *7 *7 *7 *7 *3 *7 *7 | |
| *7 *7 *7 *7 *7 *7 *7 *3 *7 *3 | 61.80 62.80 |
| B308 PILEATED WOODPECKER | 1.00 |
| DOVO TELEVIED WOODI ECKER | |
| *3 *3 | 0.00 |
| B289 CALLIOPE HUMMINGBIRD | 1.00 |
| | |
|) | 7.00 |
| 3333 WESTERN KINGBIRD | 8.10 |
| | 1.10 |
| | 43.50 |
| *10 0 0 0 *7 *3 *3 0 *3 *3 0 *7 *7 3484 CALIFORNIA TOWHEE | 44.60 |
| 97 *7 *3 *7 *7 *3 *7 *10 | 1.10 |
| | 52.90 |
| *10 *3 *7 *7 *7 *3 *10 *7 *3 *3 *7 M033 WESTERN RED BAT | 54.00 |
| | 1.10 |
| *7 | 57.80 |
| | 58.90 |
| A145 COMMON PORCUPINE | 1.10 |
| *5 *5 *5 *4 *4 *3 *3 0 | 48.60 |
| *3 *3 *5 *5 *5 *4 *4 *3 0 *7 *5 | 49.70 |
| M151 BLACK BEAR | 1.10 |
| *5 *8 *8 *5 *5 *5 *4 0 | 64.20 |
|) *3 *4 *5 *8 *8 *4 *5 *7 *8 *7 | 65.30 |
| R053 STRIPED RACER | 1.10 |
| *3 *3 *3 *7 *7 *10 | 31.80 |
| *10 *10 *3 *3 *3 *7 *7 *3 | 32.90 |
| WESTERN AQUATIC CHRIER SWARE | 1.10 |
| *5 *5 *5 *10 *10 *5 *5 *3 *10 | 51.50 |
| 3 *5 *5 *5 *5 *5 *10 *10 *5 *10 *5 *5 | 52.60 |
| 3339 TREE SWALLOW | 1.20 |
| 4 *4 *4 0 0 *4 0 0 0 *10 | 44.80 |
| 10 0 0 0 *4 *4 *4 0 0 *4 0 *7 *7 | 46.00 |
| 3369 HOUSE WREN | 1.20 |
| *10 *10 *7 *10 *10 *7 *3 *10 | 83.40 |
| *10 *3 *3 *10 *10 *7 *9 *10 *7 *10 *7 | 84.60 |
| 3411 EUROPEAN STARLING | 1.20 |
| 10 *9 *7 0 0 *4 0 0 *10 | 79.00 |
| 10 0 0 *10 *9 *7 0 0 *4 0 *10 *7 | 80.20 |
| MO27 LONG-LEGGED MYOTIS | 1.20 |
| 7 *7 *7 *7 *7 *7 0 0 0 *0 | 60.80 |
| 0 0 0 0 0 *7 *7 *7 *7 *7 *9 *7 *10 | 62.00 |
| M142 HOUSE MOUSE | 1.20 |
| 3 *3 *3 *3 *3 *10 *7 *10 | 39.00 |
| 10 *10 *7 *3 *3 *3 *7 *3 *7 *3 | 40.20 |
| 3138 WILD TURKEY | 1.30 |
| 10 *10 *9 *9 *9 *9 0 *10 | 86.70 |
| 10 0 0 *10 *10 *9 *9 *9 *10 *10 *10 | 88.00 |

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|---|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B483 SPOTTED TOWHEE | 0.80 |
| *9 *5 *4 *7 *7 *3 *10 *10 | 58.20 |
| *10 *8 *10 *9 *5 *4 *9 *7 *3 0 *7 | 59.00 |
| B495 LARK SPARROW | 0.80 |
| *7 *7 *3 *3 *3 *10 | 45.10 |
| *10 *3 *3 *7 *7 *3 *3 *4 *3 | 45.90 |
| M057 ALLEN'S CHIPMUNK | 0.80 |
| *3 *3 *3 *3 *3 *7 | 31.00 |
| 0 *7 *3 *3 *3 *3 *7 *3 *3 | 31.80 |
| B111 WHITE-TAILED KITE | 0.90 |
| *8 *7 *5 0 0 *8 0 0 *10 0 | 62.40 |
| 0 *10 0 0 *8 *8 *7 *5 0 0 0 *10 | 63.30 |
| M030 SILVER-HAIRED BAT | 0.90 |
| *7 *7 *5 0 0 0 0 0 0 | 52.70 |
| 0 0 0 0 0 *7 *7 *5 0 0 0 *7 *7 | 53.60 |
| M045 BRUSH RABBIT | 0.90 |
| *7 *5 *3 *10 *9 *9 *8 *10 | 51.00 |
| *10 *8 *8 *9 *7 *5 *3 *8 *10 *3 *3 | 51.90 |
| A001 CALIFORNIA TIGER SALAMANDER | 1.00 |
| *3 *3 *7 | 30.10 |
| *7 *3 *3 *3 *7 | 31.10 |
| B158 KILLDEER | 1.00 |
| *3 *10 *10 | 14.30 |
| *10 *10 *3 *3 *3 | 15.30 |
| B263 FLAMMULATED OWL | 1.00 |
| 0 | 0.00 |
| 0 *10 | 1.00 |
| B276 COMMON NIGHTHAWK | 1.00 |
| *5 0 0 0 0 0 0 0 *9 *10 | 16.20 |
| *10 *9 0 0 0 *5 0 0 0 0 0 *5 | 17.20 |
| B294 LEWIS' WOODPECKER | 1.00 |
| 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 *10 | 1.00 |
| B302 NUTTALL'S WOODPECKER | 1.00 |
| *10 *10 *8 0 0 0 0 *5 | 81.70 |
| <u>*5 0 0 *10 *10 *8 0 0 0 *10 *8</u> | 82.70 |
| B360 BUSHTTT | 1.00 |
| *8 *10 *10 *5 *5 *4 *7 *10 | 93.50 |
| <u>*10 *3 *7 *8 *10 *10 *5 *5 *4 *10 *7</u> | 94.50 |
| B545 AMERICAN GOLDFINCH | 1.00 |
| *5 *3 *3 *7 *5 0 0 *10 | 41.40 |
| <u>*10 0 0 0 *5 *3 *3 *10 *7 *5 0</u> | 42.40 |
| M105 CALIFORNIA KANGAROO RAT | 1.00 |
| *3 *7 | 1.00 |
| *7 *10 *3 | 2.00 |
| M120 PINON MOUSE | 1.00 |
| *7 *7 *7 *10 *5 | 3.60 |
| *5 *3 *10 *7 *7 *7 *7 | 4.60 |
| R023 SAGEBRUSH LIZARD | 1.00 |
| *3 | 0.30 |
| *3 *3 *7 | 1.30 |
| | |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|---|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B521 WESTERN MEADÓWLARK | 0.70 |
| *5 *3 *3 *3 *10 *8 | 38.80 |
| *8 *10 *3 *5 *3 *3 *5 0 | 39.50 |
| B547 HOUSE SPARROW | 0.70 |
| 0 0 0 0 *10 | 7.00 |
| *10 0 0 0 0 *3 | 7.70 |
| M025 LONG-EARED MYOTIS | 0.70 |
| *7 *7 *7 *10 *10 *7 0 0 *0 | 61.40 |
| *0 0 0 *7 *7 *7 *10 *10 *7 0 *7 *7 | 62.10 |
| M029 WESTERN SMALL-FOOTED MYOTIS | 0.70 |
| *7 *7 *3 *5 *5 *3 0 0 0 *9 *0 | 50.90 |
| *0 *9 0 0 0 *7 *7 *3 *5 *5 *3 0 *7 *3 | 51.60 |
| M038 PALLID BAT | 0.70 |
| *7 *7 *7 0 0 0 *7 0 *5 *0 | 63.10 |
| *0 *5 0 *7 *7 *7 *7 *7 0 0 0 *7 0 | 63.80 |
| M087 SAN JOAQUIN POCKET MOUSE | 0.70 |
| *10 *3 | 1.30 |
| *3 *10 *7 | 2.00 |
| M160 AMERICAN BADGER | 0.70 |
| *3 *3 *10 *7 | 8.30 |
| *7 *10 *7 *3 *3 | 9.00 |
| M165 MOUNTAIN LION | 0.70 |
| *7 *9 *9 *7 *5 *10 *3 | 76.80 |
| *3 *4 *10 *7 *9 *9 *5 *9 *9 *7 *5 | 77.50 |
| R039 WESTERN WHIPTAIL | 0.70 |
| *3 *3 *3 *3 *3 *7 *3 | 31.30 |
| *3 *3 *7 *3 *3 *3 *3 *3 *3 *3 *3 | |
| R046 RUBBER BOA | 32.00 |
| *3 *3 *10 | |
| *3 *10 *10 | 1.60 2.30 |
| B119 RED-SHOULDERED HAWK | 0.80 |
| | |
| | 7.80 |
| *10 0 0 0 0 0 *4 0 0 *6 0 B121 SWAINSON'S HAWK | 8.60 |
| | 0.80 |
| *4 *3 0 *8 *8 0 | 22.00 |
| 0 *8 *8 0 *4 *3 0 0 *5 0 | 22.80 |
| B251 BAND-TAILED PIGEON | 0.80 |
| 0 0 0 *5 0 *10 | 7.50 |
| *10 0 0 0 0 0 *5 0 *8 | 8.30 |
| B299 RED-BREASTED SAPSUCKER | 0.80 |
| 0 0 0 0 0 0 *10 | 7.00 |
| *10 0 0 0 0 0 0 0 *8 | 7.80 |
| B321 BLACK PHOEBE | 0.80 |
| *3 *3 *9 *9 *3 *3 *4 0 *10 0 | 39.20 |
| 0 *10 *4 *3 *3 *3 *3 *10 *9 *3 0 *3 *3 | 40.00 |
| B386 HERMIT THRUSH | 0.80 |
| 0 0 0 0 *10 0 0 | 1,00 |
| 0 0 0 0 0 0 0 0 *10 0 *8 | 1.80 |
| B430 YELLOW WARBLER | 0.80 |
| *6 *6 0 *10 *8 0 0 *7 | 32.40 |
| *7 0 *6 *6 0 *10 *10 0 *6 0 | 33.20 |

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|---|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| 368 BEWICK'S WREN | 0.60 |
| 9 *10 *10 *10 *9 *3 *10 *10 | 96.70 |
| 10 *7 *10 *9 *10 *10 *8 *10 *3 *9 *3 | 97.30 |
| 391 WRENTIT | 0.60 |
| 7 *3 *7 *3 *3 0 *10 *7 | 57.00 |
| 7 *9 *10 *7 *3 *7 *3 *3 0 *3 0 | 57.60 |
| 809 INDIGO BUNTING | 0.60 |
| 0 0 *3 *3 *3 *3 | 1.20 |
| 3 *3 0 0 0 *3 *3 *3 0 *3 | 1.80 |
| 077 WESTERN GRAY SQUIRREL | 0.60 |
| 7 *7 *10 0 0 *7 0 0 *7 *7 *10 0 0 *7 *8 *7 | 75.00 |
| 0 *7 *7 *10 0 0 *7 *8 *7 | 75.60 |
| 1156 ERMINE | 0.60 |
| | 0.50 |
| *7 | 1.10 |
| 004 WESTERN POND TURTLE | 0.60 |
| 0 *10 *10 *7 *7 *3 *3 *7 *10 *0 | 92.70 |
| *7 *3 *3 *10 *10 *10 *7 *7 *3 *10 *10 *3 | 93.30 |
| 003 LONG-TOED SALAMANDER | 0.70 |
| 3 *3 *7 *3 | 1.60 |
| *3 *7 *3 *7 | 2.30 |
| 007 CALIFORNIA NEWT | 0.70 |
| 10 *10 *10 *7 *7 *10 *3 *10 | 92.70 |
| 3 *7 *10 *10 *10 *10 *7 *7 *10 *10 | 93.40 |
| 012 ENSATINA | 0.70 |
| 3 *3 *3 *7 *7 *7 *3 | 31,80 |
| 3 *3 *3 *3 *3 *7 *7 *7 *3 *3 | 32.50 |
| 028 WESTERN SPADEFOOT | 0.70 |
| 3 *3 *3 *7 *8 | 16.00 |
| 7 *3 *3 *3 *3 *8 *3 | 16.70 |
| 133 RING-NECKED PHEASANT | 0.70 |
| 0 0 0 *10 *7 | 5.60 |
| *10 *3 0 0 0 *4 0 0 | 6.30 |
| 304 HAIRY WOODPECKER | 0.70 |
| *4 *3 0 0 *7 0 | 34.00 |
| 0 *4 *4 *3 0 0 *7 *3 *7 | 34.70 |
| 323 SAY'S PHOEBE | 0.70 |
| 3 0 *9 | 7.20 |
| 0 *3 *3 | 7.90 |
| 367 CANYON WREN | 0.70 |
| 7 *7 *7 *10 *10 *7 *10 | 62.40 |
| 10 *7 *7 *7 *7 *7 *10 *10 *7 | 63.10 |
| 398 CALIFORNIA THRASHER | 0.70 |
| 0 *10 *7 | 6.00 |
| **3 *10 *4 *4 0 | 6.70 |
| 171 WESTERN TANAGER | 0.70 |
| | 0.40 |
| | |
| 0 0 0 0 0 0 0 *5 0 *6 504 FOX SPARROW | 1.10 |
| | |
| 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 *7 | 0.70 |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference | æ |
|--|------------------------------------|------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values | - |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values | |
| M070 BELDING'S GROUND SQUIRREL | 0.30 | |
| *7 *3 | 1,00 | |
| *3 *7 *3 | 1.30 | |
| M085 MOUNTAIN POCKET GOPHER | 0.30 | |
| *7 *7 | 5.30 | |
| *7 *7 *3 | 5.60 | |
| M133 MONTANE VOLE | 0.30 | |
| *7 *3 | 1,00 | |
| *7 *3 *3 | 1.30 | |
| M136 LONG-TAILED VOLE | 0.30 | |
| *3 *10 *7 | 2.00 | |
| *10 *3 *3 *7 | 2.30 | |
| B496 BLACK-THROATED SPARROW | 0.30 | |
| DENOR-TIMO/TIED BITMING W | 0,00 | |
| *3 | 0.30 | |
| A043 FOOTHILL YELLOW-LEGGED FROG | 0.40 | |
| | 30.00 | |
| *3 *3 *3 0 0 *3 0 *3 *3 0 0 *3 *3 *3 0 0 *3 *3 *3 | 30.40 | |
| *3 0 0 *3 *3 *3 0 0 *3 *3 *3 B357 CHESTNUT-BACKED CHICKADEE | 0,40 | |
| | 44,80 | |
| | 45.20 | |
| *10 0 *4 *4 *4 0 0 *4 *4 *4 | 43.20 | |
| M006 ORNATE SHREW | | |
| *3 *3 *7 *7 *3 *7 *7 | 32.50 | |
| *7 *3 *3 *3 *3 *3 *7 *7 *7 *3 | 32.90 | |
| M012 TROWBRIDGE'S SHREW | | |
| *3 *3 *3 | 23.40 | |
| *3 *3 *3 *3 *3 | 23.80 | ., , |
| M134 CALIFORNIA VOLE | | |
| *3 *3 *10 *7 *7 | 6.90 | |
| *7 *10 *7 *3 *7 | 7.30 | |
| M176 WILD PIG | 0.40 | |
| *7 *10 *8 *7 *7 *8 *10 0 | 74.70 | |
| 0 *4 *10 *7 *10 *8 *3 *7 *10 *7 *7 | 75.10 | |
| B052 GREAT EGRET | 0.50 | |
| *3 *3 *3 0 0 *7 | 34,20 | |
| *7 0 *3 *3 *3 0 *5 | 34.70 | |
| B338 PURPLE MARTIN | 0.50 | |
| 0 0 0 *10 | 7.00 | |
| *10 0 0 0 0 0 *5 | 7.50 | |
| B438 HERMIT WARBLER | 0.50 | |
| 0 0 0 0 0 0 0 | 0.00 | |
| 0 0 0 0 0 0 0 0 *5 | 0.50 | |
| B536 PURPLE FINCH | 0.50 | |
| *7 *7 *7 0 *3 0 *10 | 66.00 | |
| *10 0 0 *7 *7 *7 0 0 0 *7 *8 | 66.50 | |
| M140 BLACK RAT | 0.50 | |
| *3 *3 *3 *3 *10 | 37.00 | |
| *10 *3 *3 *3 *7 *3 *3 | 37.50 | |
| A040 RED-LEGGED FROG | 0.60 | |
| *7 *7 *7 *3 *3 *3 *7 *10 | 61.60 | |
| *7 *3 *3 *7 *7 *7 *3 *3 *10 *7 *3 | 62.20 | |

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Comments | | Difference |
|--|------------------------------------|---------------|
| ID Species Name Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values | Difference |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values | |
| M042 WESTERN MASTIFF BAT | | 0.10 |
| *3 *3 *3 0 0 0 0 0 0 0 *7 | 30.10 | |
| *7 0 0 0 0 *3 *3 *3 0 0 0 0 *3 0 | 30.20 | |
| M141 NORWAY RAT | | 0.10 |
| *3 *3 *3 *7 *7 *10 | 37.80 | |
| *10 *3 *3 *3 *7 *7 *3 | 37.90 | |
| A006 ROUGH-SKINNED NEWT | | 0.20 |
| *5 *7 *7 *7 *5 *3 *3 | 60.90 | |
| *3 *5 *7 *7 *7 *7 *3 *3 *7 *3 | 61.10 | 0.20 |
| B524 BREWER'S BLACKBIRD | | 0.20 |
| *4 *5 *5 *9 *7 *5 0 *5 *10 | 54.70 | |
| *10 0 *4 *5 *5 *7 *9 *5 *5 *4 *4 | 54.90 | 0.20 |
| A014 CALIFORNIA SLENDER SALAMANDER | (2.6) | 0.30 |
| *7 *7 *10 *10 *3 *3 *3 | 63.60 63.90 | |
| *3 *3 *3 *7 *7 *7 *10 *10 *7 A044 MOUNTAIN YELLOW-LEGGED FROG | 03.90 | 0.30 |
| | 1,00 | 0.50 |
| *3 *7 | 1.30 | |
| *3 *7 *3 B051 GREAT BLUE HERON | 1,30 | 0.30 |
| *4 *4 0 0 0 *10 | 25.10 | 0.50 |
| *10 0 *4 *4 0 0 *3 *0 | 25.40 | |
| B079 MALLARD | 22.10 | 0.30 |
| *4 *4 *10 *10 *10 | 9.80 | |
| *10 *10 *7 *4 *10 | 10.10 | |
| B170 SPOTTED SANDPIPER | | 0.30 |
| 0 | 0.00 | |
| 0 *3 | 0.30 | |
| B269 BURROWING OWL | | 0.30 |
| *3 *7 *10 *10 | 9.00 | |
| *10 *10 *7 *3 *3 | 9.30 | -10. |
| B305 WHITE-HEADED WOODPECKER | | 0.30 |
| 0 | 0,00 | |
| 0 *3 | 0.30 | » |
| B309 OLIVE-SIDED FLYCATCHER | | 0.30 |
| 0 0 0 0 *3 | 0.30 | |
| *3 *3 0 0 0 0 0 0 | 0.60 | |
| B320 PACIFIC-SLOPE FLYCATCHER | | 0.30 |
| *5 *7 *7 0 0 *10 *3 *10 | 64.60 | |
| *10 *3 *5 *7 *7 0 0 *10 *5 *5 | 64.90 | 0.20 |
| B375 GOLDEN-CROWNED KINGLET | 0.00 | 0.30 |
| 0 0 0 0 | 0.00 | |
| 0 0 0 0 *3 0 | 0.30 | 0.30 |
| B493 BLACK-CHINNED SPARROW | 0.70 | 0.30 |
| *7 | 0.70 1.00 | |
| *3 *7 | 1,00 | 0.30 |
| B542 PINE SISKIN | £ 10 | 0.50 |
| 0 0 0 0 *3 0 *7 | 5.10 5.40 | |
| *7 0 0 0 0 0 0 *3 0 *3 | 5.40 | 0.30 |
| M062 LONG-EARED CHIPMUNK | 1.00 | 0.50 |
| *10 | 1.00 | |
| *3 *10 | 1.30 | |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values | Difference |
|---|--------------------------------------|------------|
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values | ***** |
| M079 DOUGLAS' SQUIRREL | Condition 2 Dain of morganica values | 0.00 |
| *3 | 0.30 | 2.30 |
| *3 0 | 0.30 | |
| M112 AMERICAN BEAVER | V.JV | 0.00 |
|) *8 | 0.80 | 0.00 |
| 0 *8 | 0.80 | |
| M139 COMMON MUSKRAT | 0.80 | 0.00 |
| *10 *10 | 3.00 | 0.00 |
| | | |
| *10 *10 *10 | 3.00 | 0.00 |
| M147 RED FOX | 0.00 | 0.00 |
| | 0.00 | |
| 0 0 0 | 0.00 | 0.00 |
| M158 AMERICAN MINK | | 0.00 |
| *10 *10 *10 | 3.00 | |
| *10 *10 *10 | 3.00 | 0.00 |
| M159 WOLVERINE | | 0.00 |
| | 0.00 | |
|) | 0.00 | |
| M163 NORTHERN RIVER OTTER | | 0.00 |
| *9 *9 0 | 1.80 | |
| 9 *9 () | 1.80 | |
| R079 GIANT GARTER SNAKE | | 0.00 |
| *10 *10 *3 *10 | 3.30 | |
| *3 *10 *10 *10 | 3.30 | |
| 3381 MOUNTAIN BLUEBIRD | | 0.00 |
| | 0.00 | |
| | 0.00 | |
| 3409 NORTHERN SHRIKE | | 0.00 |
| | 0.00 | |
|) | 0.00 | s. |
| B494 VESPER SPARROW | | 0.00 |
| | 0.00 | |
| | 0.00 | |
| 3057 CATTLE EGRET | | 0.10 |
| *3 *3 *3 *3 *0 0 0 | 30.00 | |
| 0 0 *3 *3 *3 *3 *3 0 *3 | 30.10 | |
| 3279 BLACK SWIFT | .50.10 | 0.10 |
| *3 *3 *3 0 0 0 0 0 0 *10 | 30.40 | 0.10 |
| *10 0 0 0 0 *3 *3 *3 0 0 0 *3 0 | 30.50 | |
| 3281 VAUX'S SWIFT | 30.30 | 0.10 |
| *3 *3 *3 0 0 0 0 0 *5 | 32.80 | 0.10 |
| *5 | | |
| 3342 BANK SWALLOW | 32.90 | 0.10 |
| | 2.50 | 0.10 |
| *9 *9 0 *10 0 0 *7 | 3.50 | |
| *7 0 *10 0 0 *10 *9 0 | 3.60 | 0.10 |
| M021 LITTLE BROWN MYOTIS | _ | 0.10 |
| *3 *3 *3 0 0 0 0 0 0 0 0 *0 | 29.40 | |
| *0 0 0 0 0 *3 *3 *3 0 0 0 0 *3 0 | 29.50 | |
| M036 SPOTTED BAT | | 0.10 |
| *3 *3 *3 0 0 0 0 | 29.40 | |
| 0 0 *3 *3 *3 0 0 *3 | 29.50 | |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|--|---|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values Condition 2 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values 3520 TRICOLORED BLACKBIRD | Condition 2 Sum of Weighted Values 0.00 |
| | 2.40 |
| *7 | 2.40 |
| 3522 YELLOW-HEADED BLACKBIRD | 0.00 |
| | 1.00 |
| 0 *10 0 *10 | 1.00 |
| B530 HOODED ORIOLE | 0.00 |
| *10 | 7.00 |
| *10 | 7.00 |
| B539 RED CROSSBILL | 0.00 |
| *3 | 2.30 |
| *3 | 2.30 |
| 3546 EVENING GROSBEAK | 0.00 |
| 0 0 0 0 0 0 0 0 | 0.00 |
| | 0.00 |
| B548 CLARK'S GREBE | 0.00 |
| *7 | 0.70 |
| *7 | 0.70 |
| B648 BAIRD'S SANDPIPER | 0.00 |
| 0 | 0.00 |
| 0 | 0.00 |
| B649 PECTORAL SANDPIPER | 0.00 |
| 0 | 0.00 |
| 0 | 0.00 |
| B773 AMERICAN REDSTART | 0.00 |
| 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| B798 WHITE-THROATED SPARROW | 0.00 |
| 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| B799 HARRIS'S SPARROW | 0.00 |
| 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| M003 VAGRANT SHREW | 0.00 |
| *3 *3 *10 *10 *3 *7 *10 *10 | 28.10 |
| *10 *3 *7 *3 *3 *10 *10 *3 *10 *3 | 28.10 |
| M023 YUMA MYOTIS | 0.00 |
| *7 *7 *7 0 0 0 0 0 0 *7 | 63.30 |
| *7 0 0 0 *7 *7 *7 0 0 0 0 *7 0 | 63.30 |
| M026 FRINGED MYOTIS | 0.00 |
| *8 *8 *8 0 0 0 0 0 0 *7 | 68.30 |
| *7 0 0 0 0 *8 *8 *8 0 0 0 *8 0 | 68.30 |
| M031 WESTERN PIPISTRELLE | 0.00 |
| *7 *7 *7 0 0 0 0 0 0 *7 | 59.40 |
| *7 0 0 0 0 *7 *7 *7 0 0 0 *7 0 | 59.40 |
| M039 BRAZILIAN FREE-TAILED BAT | 0.00 |
| *7 *7 *7 0 0 0 0 0 0 *5 *0 | 62.40 |
| *0 *5 0 0 0 *7 *7 *7 0 0 0 0 *7 0 | 62.40 |
| M066 YELLOW-BELLIED MARMOT | 0.00 |
| *7 | 0.70 |
| *7 | 0.70 |

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|---|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B372 MARSH WREN | 0.00 |
| 0 0 *10 | 1.00 |
| 0 0 *10 | 1.00 |
| B373 AMERICAN DIPPER | 0.00 |
| *3 *3 *10 *10 | 8.60 |
| *10 *10 *3 *3 | 8.60 |
| B376 RUBY-CROWNED KINGLET | 0.00 |
| $ \begin{array}{cccccccccccccccccccccccccccccccccccc$ | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 | 0.00 |
| | 0.00 |
| B390 VARIED THRUSH | 0.00 |
| 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| B404 AMERICAN PIPIT | 0.00 |
| 0 0 | 0.00 |
| 0 0 0 | 0.00 |
| B407 CEDAR WAXWING | 0.00 |
| 0 0 0 0 0 0 0 *10 | 7.00 |
| *10 0 0 0 0 0 0 0 0 0 | 7.00 |
| B435 YELLOW-RUMPED WARBLER | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| B437 TOWNSEND'S WARBLER | 0.00 |
| 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| B461 COMMON YELLOWTHROAT | 0.00 |
| *10 *10 *9 *10 | 3.90 |
| *9 *10 *10 *10 | 3.90 |
| B463 WILSON'S WARBLER | 0.00 |
| 0 0 0 *10 *10 0 0 0 | 2.00 |
| 0 0 0 0 0 0 *10 *10 0 0 | 2.00 |
| B491 BREWER'S SPARROW | 0.00 |
| 0 | 0.00 |
| 0 0 DAOO CAVANNALI SDADDOW | 0.00 |
| B499 SAVANNAH SPARROW | 0.00 |
| *5 *5 0 | 0.50 |
| *5 0 B506 LINCOLN'S SPARROW | 0.50 |
| 0 0 0 | 0.00 |
| | 0.00 |
| 0 0 0 0 B509 GOLDEN-CROWNED SPARROW | 0.00 |
| 0 0 0 0 0 0 0 | |
| 0 0 0 0 0 0 0 0 0 0 | 0.00 0.00 |
| B510 WHITE-CROWNED SPARROW | 0.00 |
| 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 | 0.00 |
| B519 RED-WINGED BLACKBIRD | 0.00 |
| *7 *7 *10 *10 *8 | 8.70 |
| *8 *10 *7 *7 *10 | 8.70 8.70 |
| V AV 1 IV | 0.70 |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME

Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|--|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B227 CASPIAN TERN | 0.00 |
| *3 *3 *10 | 1.60 |
| *10 *3 *3 | 1.60 |
| B233 FORSTER'S TERN | 0.00 |
| *4 *4 *0 *0 | 0.80 |
| *0 *4 *4 *0 B235 BLACK TERN | 0.80 |
| | 0.00 |
| *10 *10 | 1.00 |
| B250 ROCK DOVE | 1,00 |
| | 0.00 |
| *10 *10 | 7.00 |
| B270 SPOTTED OWL | 7.00 |
| | 0.00 |
| 0 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 B273 SHORT-EARED OWL | 0.00 |
| | |
| 0 | 2.00 |
| B291 RUFOUS HUMMINGBIRD | 2.00 |
| 0 0 0 0 0 0 0 0 0 | |
| 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| B293 BELTED KINGFISHER | 0.00 |
| *10 *10 0 *0 | 0.00 |
| *0 *10 *10 0 | 2.00 |
| B300 WILLIAMSON'S SAPSUCKER | 2.00 |
| *10 | |
| *10 | 7.00 |
| B317 HAMMOND'S FLYCATCHER | 7.00 |
| 0 0 0 *10 | 7.00 |
| *10 0 0 0 0 | 7.00 |
| B318 DUSKY FLYCATCHER | 0.00 |
| 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 | 0.00 |
| B337 HORNED LARK | 0.00 |
| *7 0 | 0.70 |
| 0 *7 | 0.70 |
| B345 GRAY JAY | 0.70 |
| *10 | 7.00 |
| *10 | 7.00 |
| B348 WESTERN SCRUB-JAY | 0.00 |
| *8 *10 *10 0 *7 *4 *7 *10 | 93.20 |
| *10 *3 *7 *8 *10 *10 0 0 *4 *10 *4 | 93.20 |
| B356 MOUNTAIN CHICKADEE | 0.00 |
| 0 0 0 0 0 0 *10 | 7.00 |
| *10 0 0 0 0 0 0 0 0 | 7.00 |
| B361 RED-BREASTED NUTHATCH | 0.00 |
| 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 | 0.00 |
| B364 BROWN CREEPER | 0.00 |
| 0 0 0 0 0 0 *10 | 7.00 |
| *10 0 0 0 0 0 0 0 | 7.00 |
| | 1.VV |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| D Species Name | Difference Conditon 1 Sum of Weighted Values |
|--|--|
| ondition 1 Habitat Suitability Values ondition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| 168 WILLET | 0.00 |
| | 0.00 |
| | 0.00 |
| 172 WHIMBREL | 0.00 |
| 0 0 | 0.00 |
| 0 0 | 0.00 |
| 173 LONG-BILLED CURLEW | 0.00 |
| 0 0 | 0.00 |
| 0 0 | 0.00 |
| 176 MARBLED GODWIT | |
| $egin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ | 0.00 0.00 |
| 183 WESTERN SANDPIPER | 0.00 |
| 0 | 0.00 |
| 0 | 0.00 |
| 185 LEAST SANDPIPER | 0.00 |
| 0 | 0.00 |
| 0 | 0.00 |
| 191 DUNLIN | 0,00 |
| | 0.00 |
| AN ANADE DIVIED DAVINGVIED | 0.00 |
| 96 SHORT-BILLED DOWITCHER | 0.00 |
| | 0.00 0.00 |
| 197 LONG-BILLED DOWITCHER | 0.00 |
| 197 EONG-BILLED DOWN CILL | 0.00 |
| | 0.00 |
| 199 COMMON SNIPE | 0.00 |
| 10 | 1.00 |
| 10 | 1.00 |
| 200 WILSON'S PHALAROPE | 0.00 |
| *8 | 0.80 |
| *8 | 0.80 |
| 213 MEW GULL | 0.00 |
| 0 | 0.00 |
| 0 214 RING-BILLED GULL | 0.00 |
| 0 0 | 0.00 |
| 0 0 | 0.00 |
| 215 CALIFORNIA GULL | 0.00 |
| *0 | 0.00 |
| 0 0 | 0.00 |
| 216 HERRING GULL | 0.00 |
| | 0.00 |
| | 0.00 |
| 217 THAYER'S GULL | 0.00 |
| | 0.00 |
| 201 OF ALICOTIC MANOCED CLEET | 0.00 |
| 221 GLAUCOUS-WINGED GULL | 0.00 |
| | 0.00 |
| | 0.00 |

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP

and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| D Species Name | Difference | ice |
|--|---|-----|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values | |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values 0.00 | |
| B115 SHARP-SHINNED HAWK | | |
| 0 0 0 0 *10 0 0 *9 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 | 7.10 7.10 | |
| | 0.00 | |
| | 0.40 | |
| 0 | 0.40 | |
| 0 0 0 0 0 0 0 *4 0 0 B124 FERRUGINOUS HAWK | 0.00 | |
| 0 0 0 0 0 0 | 0.00 | |
| | 0.00 | |
| B125 ROUGH-LEGGED HAWK | 0.00 | |
| 0 0 0 0 | 0.00 | |
| 0 0 0 0 0 | 0.00 | |
| B128 MERLIN | 0.00 | |
| 0 0 0 0 0 0 0 0 0 0 | 0.00 | |
| 0 0 0 0 0 0 0 0 0 0 0 0 0 | 0.00 | |
| B134 BLUE GROUSE | 0.00 | |
| *4 | 0.40 | |
| *4 0 0 | 0.40 | |
| B145 VIRGINIA RAIL | 0.00 | |
| 0 0 *10 | 1.00 | |
| 0 0 *10 | 1.00 | |
| B146 SORA | | |
| *10 | 1,00 1.00 | |
| *10 B148 COMMON MOORHEN | 0.00 | |
| #10 *10 | 8.00 | |
| *10 *10 | 8.00 | |
| B149 AMERICAN COOT | 0.00 | |
| *10 *10 | 8.00 | |
| *10 *10 | 8.00 | |
| B150 SANDHILL CRANE | 0.00 | |
| 0 *10 | 1.00 | |
| 0 0 *10 | 1.00 | |
| B151 BLACK-BELLIED PLOVER | 0.00 | |
| 0 | 0.00 | |
| 0 | 0.00 | |
| B156 SEMIPALMATED PLOVER | 0.00 | |
| 0 | 0.00 | |
| 0 | 0.00 | |
| B163 BLACK-NECKED STILT | 0.00 | |
| *10 | 1,00 | |
| *10 | 1.00 | |
| B164 AMERICAN AVOCET | | |
| 0 *10 | 1.00 | |
| *10 0 | 1.00 | |
| B165 GREATER YELLOWLEGS | 0.00 | |
| 0 0 0 | 0.00 | |
| 0 0 0 B166 LESSER YELLOWLEGS | 0.00 | |
| | 0.00 | |
| 0 0 0 0 | 0.00 | |

Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference |
|--|------------------------------------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values |
| B083 CINNAMON TEAL | 0.00 |
| 0 0 *3 *10 | 1.30 |
| *3 0 0 *10 B084 NORTHERN SHOVELER | 1.30 |
| | 0.00 |
| *0 *0 | 0.90 |
| *0 *9 B085 GADWALL | 0.90 |
| *7 0 | |
| *7 0 | 0.70 0.70 |
| B086 EURASIAN WIGEON | 0.70 |
| 0 0 0 | 0.00 |
| 0 0 0 | 0.00 |
| B087 AMERICAN WIGEON | 0.00 |
| *10 0 0 | 1.00 |
| 0 *10 0 | 1.00 |
| B089 CANVASBACK | 0.00 |
| *9 | 0.90 |
| *9 | 0.90 |
| B090 REDHEAD | 0.00 |
| *7 | 0.70 |
| *7 | 0.70 |
| B091 RING-NECKED DUCK | 0.70 |
| *10 | 1,00 |
| *10 | 1.00 |
| B094 LESSER SCAUP | 0.00 |
| *0 *0 | 0.00 |
| *0 *0 | 0.00 |
| B101 COMMON GOLDENEYE | 0.00 |
| 0 0 | 0.00 |
| 0 0 | 0.00 |
| B103 BUFFLEHEAD | 0.00 |
| 0 0 | 0.00 |
| 0 0 | 0.00 |
| B104 HOODED MERGANSER | 0.00 |
| 0 0 0 0 | 0.00 |
| 0 0 0 0 | 0.00 |
| B105 COMMON MERGANSER | 0.00 |
| 0 0 | 0.00 |
| 0 0 | 0.00 |
| B107 RUDDY DUCK | 0.00 |
| 0 0 *10 | 1.00 |
| 0 0 *10 | 1.00 |
| B110 OSPREY | 0.00 |
| 0 0 0 *10 *10 0 0 0 0 | 2.00 |
| 0 0 0 0 0 0 0 *10 *10 0 0 0 | 2.00 |
| B113 BALD EAGLE | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| 0 0 0 0 0 0 0 0 0 0 0 0 | 0.00 |
| B114 NORTHERN HARRIER | 0.00 |
| 0 0 0 0 *10 *10 *4 0 | 4.90 |
| 0 *4 *10 0 0 0 0 0 *10 0 | 4.90 |

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| D Species Name | | Difference |
|--|------------------------------------|------------|
| Condition 1 Habitat Suitability Values | Condition 1 Sum of Weighted Values | |
| ondition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values | 0.00 |
| 009 EARED GREBE | 4.00 | 0.00 |
| 0 | 1.00 | |
| 0 | 1.00 | 0.00 |
| 010 WESTERN GREBE | 2.72 | 0.00 |
| | 0.70 | |
| 042 AMERICAN WHITE PELICAN | 0.70 | 0.00 |
| | 0.00 | 0.00 |
| | 0.00 | |
|) 044 DOUBLE-CRESTED CORMORANT | 0.00 | 0.00 |
|) | 0.00 | 0.00 |
|) | 0.00 | |
| 049 AMERICAN BITTERN | 0.00 | 0.00 |
| | 1.00 | 0.00 |
| 0 | 1.00 | |
| 0 050 LEAST BITTERN | 1.00 | 0.00 |
| 10 LEAST BITTERN | 1.00 | 0.00 |
| 0 | 1.00 | |
| 053 SNOWY EGRET | 1.00 | 0.00 |
| 0 0 | 0.00 | 0.00 |
| *9 | 0.90 | |
| 062 WHITE-FACED IBIS | 0.90 | 0.00 |
| WIIIL-PACED IDIS | 0.70 | 0.00 |
| | 0.70 | |
| 067 TUNDRA SWAN | 0.70 | 0.00 |
| 0 | 0.00 | 0.00 |
| 0 | 0.00 | |
| 070 GREATER WHITE-FRONTED GOOSE | 0.00 | 0.00 |
| 0 | 0.00 | 0.00 |
| 0 | | -0. |
| 071 SNOW GOOSE | 0.00 | 0.00 |
| 0 | 0.00 | 0.00 |
| 0 | 0.00 | |
| 072 ROSS' GOOSE | 0.00 | 0.00 |
| 0 | 0.00 | 0.00 |
| 0 | 0.00 | |
| 075 CANADA GOOSE | 0.00 | 0.00 |
| *10 0 | 1.70 | 5.00 |
| *7 *10 | 1.70 | |
| 076 WOOD DUCK | 1.10 | 0.00 |
| 0 0 0 0 0 *8 | 5.30 | 2.00 |
| 3 0 0 0 0 0 0 0 0 | 5.30 | |
| 077 GREEN-WINGED TEAL | 2.00 | 0.00 |
| 0 *10 0 | 2.00 | 2.00 |
| *10 *10 | 2.00 | |
| 080 NORTHERN PINTAIL | 2.00 | 0.00 |
| 3 *7 0 | 1.50 | 2.00 |
| *8 *7 | 1.50 | |
| 082 BLUE-WINGED TEAL | 1.50 | 0.00 |
| *9 | 1.80 | 0.00 |
|) *9 | 1.80 | |

CALIFORNIA WILDLIFE HABITAT RELATIONSHIPS SYSTEM Supported by

CALIFORNIA INTERAGENCY WILDLIFE TASK GROUP and maintained by the

CALIFORNIA DEPARTMENT OF FISH AND GAME Database Version: 8.0

WEIGHTED HABITAT VALUE COMPARISON REPORT

| ID Species Name | Difference | <u> </u> |
|--|------------------------------------|----------|
| Condition 1 Habitat Suitability Values | Conditon 1 Sum of Weighted Values | - |
| Condition 2 Habitat Suitability Values | Condition 2 Sum of Weighted Values | |
| B116 COOPER'S HAWK | -2.00 | |
| 0 *10 *10 0 0 *7 0 0 *10 | 77.70 | |
| *10 0 0 0 0 *10 *10 0 0 *7 0 0 | 75.70 | |
| B058 GREEN HERON | -0.90 | |
| *3 *3 *3 *10 *10 *0 *9 *10 | 39.30 | |
| *10 *3 *3 *3 *10 *0 *9 *3 *0 | 38.40 | |
| B260 GREATER ROADRUNNER | -0.70 | |
| *8 *7 *5 *3 | 33.80 | |
| *3 0 *5 *8 *7 | 33.10 | |
| B370 WINTER WREN | -0.70 | |
| *3 *7 *10 | 8.00 | |
| *10 0 *3 | 7.30 | |
| M080 NORTHERN FLYING SQUIRREL | -0.60 | |
| *3 *7 *7 *3 *3 | 53.10 | |
| *3 *7 *7 *3 *3 *3 | 52.50 | |
| B259 YELLOW-BILLED CUCKOO | -0.50 | |
| *5 *5 | 1.00 | |
| 0 *5 | 0.50 | |
| B315 WILLOW FLYCATCHER | -0.40 | |
| *9 *10 | 1.90 | |
| *6 *9 | 1.50 | |
| B385 SWAINSON'S THRUSH | -0.40 | |
| 0 0 0 *3 *7 0 0 0 | 1.00 | |
| 0 0 0 0 *3 *3 0 0 0 | 0.60 | |
| B467 YELLOW-BREASTED CHAT | -0.30 | |
| *10 *10 | 2.00 | |
| *7 *10 | 1.70 | |
| B059 BLACK-CROWNED NIGHT HERON | -0.20 | |
| *3 *5 *5 *3 *3 *0 *0 *10 *5 | 48.40 | |
| <u>*5 *0 *3 *5 *5 *3 *0 *10 *5 *0</u> | 48.20 | |
| B272 LONG-EARED OWL | -0.20 | |
| *7 *9 *7 *3 *4 *4 *7 0 | 66.90 | |
| 0 0 *7 *7 *9 *7 0 *3 *4 *7 *4 | 66.70 | |
| B346 STELLER'S JAY | -0.20 | |
| 0 *4 *4 0 0 *6 *10 | 37.00 | |
| *10 0 *4 *4 0 0 *6 0 *6 | 36.80 | |
| B460 MACGILLIVRAY'S WARBLER | -0.20 | |
| 0 0 0 *7 *7 0 0 0 | 1.40 | |
| 0 0 0 0 0 *5 *7 0 0 | 1.20 | |
| B476 BLUE GROSBEAK | -0.20 | |
| *10 *10 0 | 2.00 | |
| 0 *8 *10 | 1.80 | |
| B425 ORANGE-CROWNED WARBLER | -0.10 | |
| *9 *10 *10 *10 *10 0 *7 0 | 89.20 | |
| 0 *3 *7 *9 *10 *10 *9 *10 0 *9 0 | 89.10 | |
| A042 CASCADES FROG | 0.00 | |
| *3 | 0.30 | |
| *3 | 0.30 | |
| B006 PIED-BILLED GREBE | 0.00 | |
| *10 | 1,00 | |
| *10 | 1.00 | |
| | | |